

ORIGINAL ARTICLE

Injustice at work and incidence of psychiatric morbidity: the Whitehall II study

J E Ferrie, J Head, M J Shipley, J Vahtera, M G Marmot, M Kivimäki



Occup Environ Med 2006;63:443-450. doi: 10.1136/oem.2005.022269

See end of article for authors' affiliations

Correspondence to:
Dr J E Ferrie, Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London WC1E 6BT, UK; j.ferrie@public-health.ucl.ac.uk

Accepted 24 March 2006
Published Online First 12 May 2006

Background: Previous studies of organisational justice and mental health have mostly examined women and have not examined the effect of change in justice.

Aim: To examine effects of change in the treatment of employees by supervisors (the relational component of organisational justice) on minor psychiatric morbidity, using a cohort with a large proportion of men.

Methods: Data are from the Whitehall II study, a prospective cohort of 10 308 white-collar British civil servants (3143 women and 6895 men, aged 35-55 at baseline) (Phase 1, 1985-88). Employment grade, relational justice, job demands, job control, social support at work, effort-reward imbalance, physical illness, and psychiatric morbidity were measured at baseline. Relational justice was assessed again at Phase 2 (1989-90). The outcome was cases of psychiatric morbidity by Phases 2 and 3 (1991-93) among participants case-free at baseline.

Results: In analyses adjusted for age, grade, and baseline physical illness, women and men exposed to low relational justice at Phase 1 were at higher risk of psychiatric morbidity by Phases 2 and 3. Adjustment for other psychosocial work characteristics, particularly social support and effort-reward imbalance, partially attenuated these associations. A favourable change in justice between Phase 1 and Phase 2 reduced the immediate risk (Phase 2) of psychiatric morbidity, while an adverse change increased the immediate and longer term risk (Phase 3).

Conclusion: This study shows that unfair treatment by supervisors increases risk of poor mental health. It appears that the employers' duty to ensure that employees are treated fairly at work also has benefits for health.

Most studies of relationships at work have focused either on social support or bullying. Low social support from colleagues and supervisors has been shown to be associated with increased rates of minor psychiatric morbidity and spells of psychiatric sickness absence, with effects stronger in men than women.¹⁻³ Other studies, that have examined mainly or exclusively women however, have shown that women exposed to low social support at work are at significantly higher risk of poor mental health and a decline in mental health over time.⁴⁻⁵ A considerable body of research has documented the adverse effects of bullying on health,⁶⁻⁷ in particular mental health.⁸⁻¹¹ More recently research attention has turned to the health effects of another aspect of relationships in the workplace, relational justice.¹⁻²

Relational justice forms one component of the wider concept of organisational justice, which in addition to the interactional or relational component is comprised of a procedural component and a distributive component. The procedural component refers to the perceived fairness of formal decision making procedures, the distributive component refers to the perceived fairness of the decisions themselves, while the relational component refers to the fairness with which employees perceive they are treated by their supervisors.¹³ Previous studies examining the association between justice and health have shown relational justice to be associated cross-sectionally and prospectively with poor self-rated health, increased risk of minor psychiatric morbidity, and more medically certified absence.¹²⁻¹⁶ These studies have all been carried out among Finnish municipal workers of whom the great majority are women. However, one small study, limited to hospital physicians, showed that low relational justice was associated with an increased risk of psychological distress among men but not women.¹⁷

Very little research has examined the effects of change in relational justice on health. In the only published study to date, we used data from the Whitehall II study, a cohort of British civil servants, to show that women and men exposed to low relational justice or an adverse change in relational justice were at increased risk of poor self-rated health over five years later. Among men these associations were independent of job demands, control at work, work social support, and effort-reward imbalance, but this was not the case among women.¹⁸

Two main questions emerge from the existing literature on relational justice. Firstly, is an adverse change in relational justice associated with an increase in poor mental health (minor psychiatric morbidity) in the same way that it is related to an increase in poor self-rated general health? Secondly, do the associations between relational justice and health generally observed in women apply to men?

Although both the self-rating of health and the estimation of minor psychiatric morbidity using the General Health Questionnaire (GHQ) involve a cognitive appraisal process, they measure different health domains. While self-rated health is a measure of general health, which has been shown to be strongly associated with physical ill-health,¹⁹⁻²² the GHQ is an instrument designed to detect "cases" as opposed to "non-cases" of minor psychiatric morbidity in both clinical and non-clinical populations.²³ It is, therefore, a dimension specific instrument that allows a basic distinction to be made between those who are experiencing some kind of psychiatric disturbance and those who are not. Previous work has also shown that associations between other psychosocial characteristics of the work environment and self-rated health and minor psychiatric morbidity differ in a number of important ways.¹⁶⁻²⁴⁻²⁶

The prevalence of mental ill-health, including minor psychiatric morbidity, is not equally distributed between the sexes; the prevalence being greater among women than men. However, although minor psychiatric morbidity is a stronger predictor of short spells of psychiatric sickness absence in women, it is a stronger predictor of long spells of absence in men.² Reflecting this finding, work from the Renfrew and Paisley study has shown that the long term suicide risk associated with minor psychiatric morbidity is much stronger in men than women.²⁷ Minor psychiatric morbidity has also been shown to be consistently associated with an increased risk of coronary heart disease in men, but not in women in both the Whitehall II and Renfrew and Paisley studies,^{28, 29} and analyses of population data from the Netherlands (the NEMESIS study) concluded that mental disorders are a more important risk factor for sickness absence among men than among women.³⁰

In this study we examine associations of relational justice and change in relational justice with mental health, measured as minor psychiatric morbidity, using longitudinal data from the Whitehall II cohort of British civil servants, over two thirds of whom are men.

METHODS

The target population for the Whitehall II study was all London based office staff working in 20 civil service departments between 1985 and 1988 and aged 35–55 on entry to the study. With a response rate of 73%, the final cohort consisted of 10 308 participants (3413 women and 6895 men).³¹ The true response rate was higher, however, because around 4% of those invited were not eligible for inclusion. Although mostly white-collar, respondents covered a wide range of grades from office support to permanent secretary, the highest grade in the British Civil Service.

Baseline screening (Phase 1) took place between late 1985 and early 1988. This involved a clinical examination in which height, weight, blood pressure, and serum cholesterol were determined, among other anthropometric and biomedical measures. A self-administered questionnaire containing sections on demographic characteristics, health, lifestyle factors, work characteristics, social support, life events, and chronic difficulties was completed by each respondent at home and checked for completion at the clinic. Non-responders to the initial invitation, which was delivered to potential participants at their place of work, were followed by up to two reminder letters. From Phase 1 onwards participants could indicate whether they wanted to be contacted at work or at home, and non-responders were followed up by telephone as well as by letter. In 1989/90 (Phase 2, response rate 79% of participants at Phase 1), the same questionnaire data were collected by post. The Phase 3 (1992–93, response rate 81% of participants at Phase 1) data collection included a repeat of the clinical examination in addition to a questionnaire. Full details of all contact procedures and all methods are contained in the Whitehall II Phase 3 manual.³²

Measures

Relational justice

Organisational justice has not been measured directly in the Whitehall II study. However, it has been possible to construct an indicator of *relational justice* with face validity from questions available in the Phase 1 and 2 surveys. Of the 19 items covering management and organisation of work, five items that deal with relational justice were identified using factor analysis.

- Do you get consistent information from line management (your superior)?

Table 1 Descriptive statistics for participants who were not GHQ cases at baseline and for whom data were available for relational justice at Phase 1 and GHQ at Phase 2 or Phase 3

Characteristics	Women		Men		p value for sex difference in characteristics	
	n (%)	Mean (SE)	Relational justice score (SE)	n (%)		Mean (SE)
Phase 1						
Relational justice	1975		80.1 (0.3)	4641	80.1 (0.2)	0.53
Age group (years)						<0.001
35–39	441 (22)		80.0 (0.6)	1315 (28)	80.2 (0.3)	
40–44	435 (22)		80.2 (0.6)	1265 (27)	79.9 (0.3)	
45–49	461 (23)		80.2 (0.6)	906 (20)	80.0 (0.4)	
50–55	638 (32)		80.8 (0.5)	1155 (25)	80.4 (0.3)	
Grade						<0.001
Administrative	215 (11)		80.9 (0.9)	1829 (39)	81.1 (0.3)	
Professional/executive	762 (39)		80.9 (0.4)	2421 (52)	77.3 (0.6)	
Clerical/support	998 (51)		79.4 (0.5)	391 (8)	79.8 (0.2)	
Physical illness indicator‡						0.34
No	1770 (90)		80.4 (0.3)	4195 (90)	80.2 (0.2)	
Yes	205 (10)		79.7 (0.9)	446 (10)	79.2 (0.6)	
Job demands	1957	50.8 (0.4)		4632	58.9 (0.3)	<0.001
Job control	1943	58.0 (0.3)		4620	69.4 (0.2)	<0.001
Social support at work	1956	75.4 (0.4)		4632	77.7 (0.3)	<0.001
Effort–reward imbalance	1972	0.96 (0.005)		4640	1.03 (0.003)	<0.001
Phase 2						
Relational justice	1719		78.8 (0.3)	4047	78.5 (0.2)	0.47
GHQ case						<0.001
No	1317 (76)		80.9 (0.4)	3275 (80)	80.8 (0.2)	
Yes	418 (24)		78.0 (0.6)	800 (20)	78.2 (0.4)	
Phase 3						
GHQ case†						<0.001
No	997 (66)		81.3 (0.4)	2650 (13)	80.9 (0.2)	
Yes	505 (34)		78.1 (0.6)	1020 (27)	78.8 (0.4)	

†GHQ case at Phase 2 or Phase 3.

‡Diabetes, diagnosed heart trouble, hypertension, ECG abnormalities, and/or respiratory illness.

- Do you get sufficient information from line management (your superior)?
- When you are having difficulties at work, how often is your superior willing to listen to your problems?
- Do you ever get criticised unfairly? (reverse scored)
- Do you ever get praised for your work?

These five items formed an internally consistent scale of relational justice (Cronbach's alpha 0.71 at Phase 1, 0.73 at Phase 2). Responses were scored on a four point scale from never (1) to often (4). The mean of scores were scaled from 25 to 100. For example, a participant who selected "often" in answer to every question would score 100 while a participant who selected "never" would score 25. The resulting distribution was divided into tertiles (25–70 = "low", 71–88 = "intermediate", 89–100 = "high"). The bottom tertile indicated a low level of relational justice, the top tertile a high level of relational justice, and the middle tertile an intermediate level. Change in relational justice was calculated by deducting the Phase 1 score from the Phase 2 score. Participants were classified into one of three groups: "no change", "adverse change" (a decline of 10 score points or more), and "favourable change" (an increase of 10 score points or more).

Other psychosocial characteristics of the work environment

The demand-control and the effort-reward imbalance model represent the two main models of organisational stress. Subsequent to its initial conception, social support was added to the demand-control model. Forty self-report questions based on the constructs of the demand-control-support model, derived from well known questionnaires used in the United States and Sweden and the Job Content Questionnaire, were included in the Phase 1 questionnaire.³³⁻³⁵ Reliability analysis and exploratory principal components analysis at Phase 1 showed that some of the items were not reliable measures of the underlying construct.³⁶ Furthermore, some items were deleted from the scale as they were left unanswered by too many participants. For such reasons 15 of the 40 questions were dropped from successive questionnaires. Of the remaining 25 questions, 15 measure *job control* or decision authority, 9 of the 15 items cover decision latitude, and 6 cover skill discretion. These subscales were equally weighted (internal consistency, Cronbach's $\alpha = 0.84$). *Job demands* were measured using 4 items (internal consistency, Cronbach's $\alpha = 0.67$), and *social support* at work, which comprised three components (support from colleagues, support from supervisors, and clarity and consistency of information from supervisors), was measured using 6 items (internal consistency, Cronbach's $\alpha = 0.79$). A list of all 25 questions has been published previously.³⁷ All questions were answered on a four point scale from "often" to "never/almost never". Responses on a four point scale from "often" to "never/almost never" were combined into summary scales and then divided into tertiles. In all measures, the bottom tertile indicated a low level, the top tertile a high level, and the middle tertile an intermediate level for each of these indicators.³⁸ The *effort-reward* model of work stress in its current form was not available at Phase 1, so a measure derived from existing questions was used. Details of the derivation and testing of this measure have been published previously.³⁹

Minor psychiatric morbidity

The 30 item General Health Questionnaire (GHQ) was used to assess psychiatric morbidity.²³ GHQ items consist of statements about behavioural and psychological functioning. The respondent is asked to say how well the statement applies to them during "the past few weeks" in comparison

Table 2 Associations between relational justice at Phase 1 and GHQ caseness by Phase 2

		Covariates at Phase 1					All
		Unadjusted	Age, grade, physical illness†	Age, grade, physical illness, and job demands	Age, grade, physical illness, and job control	Age, grade, physical illness, and social support	Age, grade, physical illness, and effort-reward imbalance
n (cases)		OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Women*	1 688 (406)						
Relational justice							
High	578 (118)	1.00	1.00	1.00	1.00	1.00	1.00
Intermediate	454 (101)	1.12 (0.82 to 1.51)	1.06 (0.78 to 1.44)	1.05 (0.78 to 1.43)	1.04 (0.77 to 1.41)	1.07 (0.79 to 1.45)	1.06 (0.77 to 1.47)
Low	656 (187)	1.55 (1.19 to 2.02)	1.51 (1.16 to 1.98)	1.47 (1.12 to 1.93)	1.45 (1.11 to 1.91)	1.51 (1.13 to 2.00)	1.43 (0.99 to 2.06)
Test for linear trend		p < 0.0001	p < 0.001	p < 0.001	p < 0.001	p < 0.001	p = 0.02
Men*	4048 (797)						
Relational justice							
High	1234 (200)	1.00	1.00	1.00	1.00	1.00	1.00
Intermediate	1370 (255)	1.18 (0.96 to 1.44)	1.16 (0.94 to 1.42)	1.15 (0.94 to 1.41)	1.13 (0.92 to 1.39)	1.14 (0.92 to 1.43)	1.07 (0.87 to 1.32)
Low	1444 (342)	1.60 (1.32 to 1.95)	1.63 (1.34 to 1.98)	1.59 (1.31 to 1.93)	1.54 (1.26 to 1.88)	1.53 (1.18 to 1.99)	1.37 (1.11 to 1.68)
Test for linear trend		p < 0.0001	p < 0.001	p < 0.001	p < 0.001	p < 0.001	p = 0.001

*Excludes GHQ cases at baseline and participants with missing data in any of the predictors.
†Diabetes, diagnosed heart trouble, hypertension, ECG abnormalities, and/or respiratory illness.

to their “usual” behaviour or state of mind. For example: “Have you recently felt you are playing a useful part in things?”, “Have you recently felt you couldn’t overcome your difficulties?”. The response alternatives are: “not at all”, the “same as usual”, “rather more than usual”, and “much more than usual”. As little is gained in terms of case identification by discriminating between the severity of symptoms, questions were scored as follows: “not at all” = 0, “same as usual” = 0, “rather more than usual” = 1, and “much more than usual” = 1. The GHQ-30 has been validated in a number of diverse populations and has been validated specifically against the Clinical Interview Schedule in Whitehall II data, giving a cut-off point of 4/5 for dividing “non-cases” from “cases”. GHQ caseness was thus defined as a score of 5 or more. Cases are participants who are at higher risk of minor psychiatric morbidity, largely depression, and anxiety disorders.^{24–40}

Covariates

Age and employment grade were derived from the Phase 1 questionnaire. Employment grade was determined from the participant’s Civil Service grade title. For analysis, employment grade titles were divided into three categories in order of decreasing salary: administrative, professional/executive, and clerical/support. A composite indicator of physical illness (*physical illness indicator*) was comprised of diabetes, diagnosed heart trouble, ECG abnormalities, hypertension, and/or respiratory illness. The category “diabetes” included all diabetics. Data on past medical history of doctor diagnosed coronary heart disease (CHD) were derived from the Phase 1 questionnaire. ECG abnormalities were probable/possible ischaemia identified on ECG during the baseline screening examination. The category “hypertension” included all participants on antihypertensive medication or with a systolic or diastolic blood pressure greater than 160 or 95 mm Hg respectively. Presence of a respiratory illness was detected using the Medical Research Council chronic bronchitis questionnaire.⁴¹

Statistical analysis

Associations between relational justice at Phase 1 and GHQ caseness by Phases 2 and 3 were determined using logistic regression analyses among participants who were not GHQ cases at baseline. The outcome of interest in these analyses at Phase 2 was prevalence of GHQ caseness by Phase 2, while at Phase 3 it was GHQ caseness by Phase 3 (GHQ case at Phase 2 or at Phase 3). Results are presented as odds ratios and 95% confidence intervals adjusted for age in five year categories, employment grade, and physical illness at Phase 1. The physical illness indicator was included as a covariate in order to control for physical illness and reduce the effects of reverse causality. A linear trend between relational justice and GHQ caseness was tested by entering the continuous justice score in the model.

Three of the questions used to construct our relational justice measure are from the social support scale and one from the effort–reward imbalance model. In order to determine whether our measure of relational justice had explanatory power beyond that provided by social support at work and the existing well known work stress models, the next step in our analyses was to adjust for these work characteristics, firstly separately and then all together. The effect of these adjustments was summarised by calculating the percentage change in the linear trend term for relational justice. Only participants with no missing data for any of the work characteristics were included in these models.

The next step in these analyses tested whether favourable or adverse change in relational justice between Phase 1 and Phase 2, compared with no change, predicted GHQ caseness

by Phase 2 and Phase 3 after adjustment for age, employment grade, physical illness, and relational justice at Phase 1. Linear trend was tested by entering the change score in the model. All analyses were conducted separately for women and men using the SAS statistical program (SAS Institute, Cary, NC, USA).

Ethical approval for the Whitehall II study was obtained from the University College London Medical School Committee on the ethics of human research.

RESULTS

At Phase 1, 2744 participants had psychiatric disorder as indicated by GHQ caseness, and 119 had missing data on the GHQ. Both these groups were excluded from the main analyses. Of the 7445 participants free of psychiatric disorder at Phase 1 (non-GHQ cases), 7434 responded to justice items at Phase 1, 5810 also responded to the GHQ at Phase 2, and 5172 additionally responded to the GHQ at Phase 3 (table 1). In the analyses of justice at Phase 1 and GHQ incidence by Phase 2, those 5736 participants who additionally had no missing data in any of the Phase 1 covariates were included (table 2). In the analyses of GHQ caseness by Phase 3, the number of included participants was 5109 (table 3). Analyses of change in justice between Phase 1 and Phase 2 were adjusted for age, grade, physical illness, and relational justice at Phase 1 (but not for the other baseline covariates); among these, 5700 participants had complete data for incident GHQ at Phase 2 and 5086 for incident GHQ by Phase 3 (table 4). The latter population (the group with the greatest attrition) compared with all non-GHQ cases at Phase 1 contained a slightly smaller proportion of women (29% v 31%) and manual workers (19% v 23%), but any differences in the prevalence of physical illness (9% v 10%) and the level of justice (80.3 v 80.1), job demands (56.5 v 56.0), job control (66.4 v 65.5), social support at work (77.4 v 76.7), and effort–reward imbalance (1.0 v 1.0) were minimal or non-existent. Participants who were GHQ cases at baseline had lower relational justice score (75.4) than included participants.

As can be seen in table 1, in contrast to the other psychosocial characteristics of the work environment, women and men reported the same mean levels of organisational justice at Phase 1. Furthermore, there were few differences in relational justice scores by age or by grade, with the exception of men in the middle grades who had a slightly lower relational justice score than men in the highest and lowest grades. However, relational justice scores for both sexes were lower at Phase 2 than at baseline but again did not differ between women and men. While statistically significant ($p < 0.001$), this decline in relational justice was relatively small (0.12 SD).

Tables 2 and 3 show strong associations ($p < 0.001$) between relational justice at Phase 1 and GHQ caseness by Phases 2 and 3 in both sexes. Among women the association seems to be driven mainly by the significantly higher risk of GHQ caseness associated with low relational justice. Similar results are seen in men, apart from the intermediate relational justice group in which there appears to be a slightly higher risk of GHQ caseness by both phases. With regard to low relational justice, adjustment for age, grade, and the physical illness indicator produces figures little different from those seen in the unadjusted associations.

The association between relational justice and GHQ caseness in women is attenuated by 12% for caseness by Phase 2 and 9% for caseness by Phase 3 when job control is included in the model adjusted for age, grade, and the physical illness indicator. Social support produces little attenuation of the association at Phase 2, but the association with caseness by Phase 3 is attenuated by 30%. When all the other characteristics of the work environment are added into the

Table 3 Associations between relational justice at Phase 1 and GHQ caseness by Phase 3 (GHQ cases at Phase 2 or Phase 3)

		Covariates at Phase 1						
		Unadjusted	Age, grade, physical illness†	Age, grade, physical illness, and job demands	Age, grade, physical illness, and job control	Age, grade, physical illness, and social support	Age, grade, physical illness, and effort-reward imbalance	All
n (cases)		OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Women*	1465 (496)							
Relational justice								
High	498 (145)	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Intermediate	397 (118)	1.03 (0.77 to 1.38)	0.97 (0.72 to 1.30)	0.97 (0.73 to 1.31)	0.95 (0.71 to 1.28)	0.94 (0.69 to 1.29)	0.98 (0.73 to 1.32)	0.94 (0.68 to 1.28)
Low	570 (233)	1.68 (1.30 to 2.17)	1.64 (1.27 to 2.13)	1.62 (1.24 to 2.10)	1.58 (1.21 to 2.05)	1.45 (1.03 to 2.06)	1.64 (1.24 to 2.16)	1.40 (0.98 to 2.00)
Test for linear trend		p < 0.0001	p < 0.001	p < 0.001	p < 0.001	p = 0.007	p < 0.001	p = 0.03
Men*	3644 (1012)							
Relational justice								
High	1114 (265)	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Intermediate	1228 (333)	1.19 (0.99 to 1.44)	1.17 (0.97 to 1.41)	1.15 (0.95 to 1.39)	1.15 (0.95 to 1.39)	1.20 (0.98 to 1.47)	1.10 (0.91 to 1.34)	1.14 (0.93 to 1.40)
Low	1302 (414)	1.49 (1.25 to 1.79)	1.52 (1.27 to 1.82)	1.46 (1.22 to 1.76)	1.47 (1.22 to 1.77)	1.59 (1.24 to 2.03)	1.32 (1.09 to 1.60)	1.43 (1.11 to 1.84)
Test for linear trend		p < 0.0001	p < 0.001	p < 0.001	p < 0.001	p < 0.001	p < 0.001	p < 0.001

*Excludes GHQ cases at baseline and participants with missing data in any of the predictors.
 †Diabetes, diagnosed heart trouble, hypertension, ECG abnormalities, and/or respiratory illness.

model together, the association between relational justice and GHQ caseness in women is attenuated by 16% at Phase 2 and 38% for caseness by Phase 3. Among men, job demands, job control, and social support produce a modest attenuation (6–16%) of the association between relational justice and GHQ caseness by Phase 2 and by Phase 3, whereas effort–reward imbalance attenuates the association by 41% at Phase 2 and by 38% for caseness by Phase 3. At Phase 2 the greatest attenuation, 46%, is seen when all the work characteristics are entered into the model together. However, the test for trend remains statistically significant for the fully adjusted model across both phases and sexes.

Table 4 shows that in both sexes a favourable change in relational justice between Phase 1 and Phase 2 is associated with a decreased risk of incident GHQ caseness at Phase 2 compared with the group that experienced no change. Conversely, an adverse change is associated with an increased risk. In both sexes the effect of a favourable change in relational justice between Phase 1 and Phase 2 on GHQ caseness appears to have weakened by Phase 3. However, the effects of an adverse change remain strong by Phase 3 and the test for linear trend between change in justice and GHQ caseness remains unchanged at both phases (p < 0.001).

DISCUSSION

Women and men who reported at baseline that they were treated unfairly by their supervisors had a significantly higher risk of incident psychiatric morbidity three years and six years later. Less than half of this effect appeared to be explained by other psychosocial characteristics of the work environment, the most important explanatory factor being social support at work among women and effort–reward imbalance among men. Overall there was a small decline in the level of relational justice among Whitehall II participants of both sexes over the three years following the baseline survey. Within this overall decline, some participants experienced a favourable change in justice, while others experienced an adverse change. A favourable change was associated with a significant reduction in the immediate risk of psychiatric morbidity, while an adverse change significantly increased both the immediate risk and the longer term risk.

This study benefits from using data from the Whitehall II study, a well characterised cohort with sufficient power to detect effects in both sexes. It is also, to our knowledge, the first study to examine the effects of change in relational justice on mental health in the workplace. By analysing change in a sub-population comprised of participants who were not GHQ cases at baseline, the study addressed the possibility of reverse causation, that is the situation in which participants with higher GHQ scores are treated less fairly by supervisors, rather than GHQ caseness being the result of such unfair treatment. In the absence of controlled experiments, the examination of change in an exposure over time, while removing prevalent cases of the outcome at baseline, provides the best means of tackling such reverse causation in addition to other problems, such as bias and confounding, which can be controlled in the experimental situation. Adjusting for a measure of physical illness at baseline also reduced the likelihood that our findings could be attributable to pre-existing physical morbidity.

Although using a sub-population of non-GHQ cases at baseline reduces the possibility of reverse causation, it remains possible that borderline GHQ caseness also elicited less fair treatment. To control for the effects of sub-clinical psychiatric morbidity and to determine the effects of relational justice and change in relational justice in the whole study population, we repeated the analyses in a

Table 4 Change in relational justice between Phase 1 and Phase 2 as a predictor of GHQ caseness by Phase 2 or by Phase 3

Change between Phase 1 and Phase 2	GHQ caseness by Phase 2			GHQ caseness by Phase 3		
	n (cases)	Unadjusted OR (95% CI)	Adjusted* OR (95% CI)	n (cases)	Unadjusted OR (95% CI)	Adjusted* OR (95% CI)
Women	1684 (404)			1464 (491)		
Relational justice						
Favourable change	383 (82)	0.94 (0.71 to 1.26)	0.74 (0.55 to 1.01)	331 (114)	1.18 (0.90 to 1.55)	0.94 (0.70 to 1.26)
No change	964 (216)	1.00	1.00	844 (260)	1.00	1.00
Adverse change	337 (106)	1.59 (1.21 to 2.09)	1.74 (1.31 to 2.30)	289 (117)	1.53 (1.16 to 2.01)	1.68 (1.26 to 2.24)
Test for linear trend		p < 0.0001	p < 0.0001		p = 0.02	p < 0.0001
Men	4016 (785)			3622 (1008)		
Relational justice						
Favourable change	808 (144)	0.98 (0.79 to 1.20)	0.75 (0.60 to 0.94)	728 (199)	1.06 (0.87 to 1.27)	0.86 (0.70 to 1.05)
No change	2471 (449)	1.00	1.00	2246 (590)	1.00	1.00
Adverse change	737 (192)	1.59 (1.31 to 1.93)	1.81 (1.48 to 2.21)	648 (219)	1.43 (1.19 to 1.73)	1.59 (1.31 to 1.93)
Test for linear trend		p < 0.0001	p < 0.0001		p = 0.0003	p < 0.0001

*Adjusted for age, grade, and physical illness indicator (diabetes, diagnosed heart trouble, hypertension, ECG abnormalities, and/or respiratory illness) and relational justice at Phase 1.

population that included all participants irrespective of GHQ status at baseline and excluded only the 1247 with missing data. In addition to adjustment for age, grade, and the physical illness indicator at baseline, these analyses were also adjusted for baseline GHQ score. The outcome of interest in these analyses at Phase 2 was prevalence of GHQ caseness at Phase 2, while at Phase 3 it was GHQ caseness at both Phase 2 and Phase 3, an outcome chosen to capture the effect of relational justice on longer term psychiatric morbidity. On adjustment for GHQ score at baseline, in general, these analyses produced findings little different from those presented (tables available from the authors on request). The two main differences were that the association between low relational justice and caseness among women at Phase 2 was attenuated by 32% rather than 16% on adjustment for all the work characteristics, and the test for trend for the change score and GHQ caseness at Phase 3 in women was $p = 0.02$ instead of $p < 0.001$. Similarly, although GHQ caseness at both Phase 2 and Phase 3 among the total cohort is not strictly comparable to the outcomes used in the main analyses, the findings followed the same pattern seen for GHQ caseness by Phase 3 (GHQ case at Phase 2 or at Phase 3) among participants case-free at baseline. The only deviation was that the association between low relational justice and GHQ caseness among all men at Phase 3 was attenuated by 63% instead of 33% on adjustment for all the work characteristics.

In the present study findings were very similar in both sexes. However, while low relational justice at baseline was a significant predictor of psychiatric morbidity at follow up in both sexes, there was an indication that both low and intermediate relational justice were predictive in men. These findings reflect those of previous work on relational justice. Studies have demonstrated cross-sectional associations with prevalent psychiatric morbidity,¹² and longitudinal associations in both sexes between low relational justice at baseline and psychiatric morbidity at follow up, two years later,¹⁴ although relational justice appears not to predict doctor diagnosed psychiatric disorders in women.⁴² The findings also follow a similar pattern to those seen between relational justice and self-rated health in the Whitehall II cohort, although the size of the effect of low relational justice and adverse change in relational justice on GHQ caseness is greater than the effect on self-rated health.¹⁸ Given the self-appraisal element involved in both measures one might expect to see some overlap in the findings, but it is important to stress that self-rated health and the GHQ measure different health domains.

With the exception of social support in women and effort-reward imbalance in men, only a modest fraction of the association between relational justice and psychiatric morbidity appeared to be explained by any other psychosocial work characteristic measured in the present study. The finding for social support at work among women is unsurprising given the considerable overlap between our relational justice and social support scales. Previous work that has examined whether associations between relational justice and psychiatric morbidity are independent of other psychosocial work characteristics has reported mixed findings. In results similar to those of the present study, the cross-sectional association between relational justice and psychiatric morbidity was attenuated, but survived, adjustment for job control and social support at work in a cohort of Finnish hospital personnel, mostly women.¹² However, later findings from the same study found that the prospective association between relational justice and psychiatric morbidity did not survive adjustment for decision authority.

No previous study appears to have examined the effect of adjusting the relational justice-health association for effort-reward imbalance. The effort-reward imbalance model is based on the concept of reciprocity, where effort at work is reciprocated by rewards that include salary, esteem, promotion opportunities, and job security. An imbalance between efforts and rewards has been shown adversely to affect self-esteem and predict a range of illnesses in employees.^{39 43 44} One of the items comprising our relational justice scale comes from the reward construct of the effort-reward model and an examination of the two constructs indicate that both are likely to have similar effects on self-esteem.⁴⁵ In addition to being a measure of stress at work, the effort-reward concept could also be viewed as a tolerable proxy for distributive justice. As distributive justice relates to the perceived fairness of formal decision making procedures, this is likely to be highly correlated with perceptions of just treatment by supervisors. Effort-reward imbalance has been examined prospectively in relation to GHQ caseness in Whitehall II and high effort and low reward separately and in combination were found to be predictive of GHQ caseness in men, but not in women.³

In the new labour market, characterised by increasing competition, downsizing, and tough, "macho" management styles, workplace bullying, recently described as "the silent epidemic", appears to be on the increase.⁴⁶ Although undocumented as yet, organisational justice is also likely to prove a casualty under such circumstances. During the baseline survey of the Whitehall II study the jobs of British

Main messages

- Employees who perceive that they are treated unfairly by their supervisors are at increased risk of poor mental health.
- An increase in unfair treatment increases the risk of poor mental health.
- An increase in fair treatment reduces the risk of poor mental health.
- Effects are only partially explained by other established occupational stressors.

civil servants were still viewed as jobs for life. However, by the time of the Phase 2 data collection, rumours of the future privatisation of Civil Service functions were starting to circulate,⁴⁷ providing a possible explanation for the overall decline in perceived relational justice between Phases 1 and 2. In such a situation fairness is important to people, because it helps them to deal with uncertainty,⁴⁸ in that solid, firmly constructed fairness judgments can help remove uncertainty or at least alleviate much of the discomfort that it would otherwise generate.

Study limitations

The main limitation of this study is the absence of standard measures of organisational justice. Other limitations include loss to follow up, the lack of an externally assessed measure of mental health, and inability to include work group level measures into the analyses.

Standard measures of organisational justice were not available at the time of the baseline survey of the Whitehall II study. Using existing questions we were able to derive a five item measure of relational justice. The Cronbach's alphas for our relational justice scale at Phases 1 and 3 were 0.71 and 0.73 respectively. A Cronbach's alpha of 0.70 is seen as acceptable,⁴⁹ and this is particularly true for a short measure assessing a relatively broad concept, as in this study. Although three of the items were those that also form the work social support scale and one that also forms the effort–reward imbalance scale, in both women and men the association between low relational justice and psychiatric morbidity was only partially attenuated by adjustment for social support at work and effort–reward imbalance. This indicates that our measure of relational justice has effects that are independent of both measures. Although we feel that effort–reward imbalance represents a tolerable proxy for distributive justice, there is nothing in the Whitehall II dataset that could be used to construct a measure of procedural justice, the third component of organisational justice. This is unfortunate because procedural justice has been shown to be at least as important a determinant of health as relational justice.^{14 16}

The main analyses, those presented in tables 2–4, dealt with participants free of psychiatric disorders at baseline, and of this population up to 33% were lost to follow up due to missing data. However, any differences in baseline characteristics between included and excluded civil servants were small, suggesting that a major bias is unlikely.

Although the GHQ is not an externally assessed measure of mental health, it is widely used, has been validated in a number diverse populations, and has been validated specifically against the Clinical Interview Schedule in Whitehall II cohort.²⁴ Individual perceptions of justice are not independent of the experiences of other members of the work group. However, we were unable to take into account the effect of work group in the present study. Members of the work group

Policy implications

- Results indicate that the duty of employers to ensure employees are treated fairly at work also has benefits for mental health and wellbeing.

can be important sources of justice evaluations, because they work together, and share opinions and emotions with each other.⁵⁰ Thus, the characteristics and opinions of other members of the work group may have an impact on attitudes and behaviours of the individual employee. Only a few studies have been able to test the effect of the work group. These have reported that employees within the same group share common perceptions of justice in the organisation, as indicated by a significant variation in justice perceptions both within and between work groups.^{51–53}

Implications

In addition to effects on mental health, failure to ensure fair treatment may lead to low morale, reduced work output and quality of service, and lost resources—because people who are trained and experienced leave the organisation.⁵⁴ While the increased risk of psychiatric morbidity associated with an adverse change in relational justice gives cause for concern, the fact that a favourable change decreases the risk gives cause for optimism. Interpersonal relationships between supervisors and employees in most large organisations are governed by internal rules and procedures, and some aspects are subject to statutory requirements. It is also possible to train supervisory staff in interpersonal skills and good practice. These findings indicate that in addition to reducing adverse organisational behaviours, such as intention to quit, the application of such measures may also reduce the burden of mental ill-health in the workplace.

Conclusion

The main findings from this study show that employees who perceive that they are treated unfairly by their supervisors are at increased risk of poor mental health. Moreover, a decline in perceived fair treatment increases the risk, whereas an increase in perceived fair treatment decreases the risk. These findings give further support to the notion of a health damaging or health promoting psychosocial work environment and demonstrate that the duty of employers to ensure employees are treated fairly at work may also have benefits for health.

ACKNOWLEDGEMENTS

We thank all participating Civil Service departments and their welfare, personnel, and establishment officers; the Occupational Health and Safety Agency; the Council of Civil Service Unions; all participating civil servants in the Whitehall II study; and all members of the Whitehall II study team.

Authors' affiliations

J E Ferrie, J Head, M J Shipley, M G Marmot, International Centre for Health and Society, Department of Epidemiology and Public Health, University College London Medical School, UK

M Kivimäki, J Vahtera, Finnish Institute of Occupational Health and University of Helsinki, Finland

Funding: The Whitehall II study has been supported by grants from the Medical Research Council; British Heart Foundation; Health and Safety Executive; Department of Health; National Heart Lung and Blood Institute (HL36310), US, NIH: National Institute on Aging (AG13196), US, NIH; Agency for Health Care Policy Research (HS06516); and the John D and Catherine T MacArthur Foundation Research Networks on Successful Midlife Development and Socio-economic Status and Health. MK, also

working at the University of Helsinki, Finland, and JV were supported by the Academy of Finland (projects 77560, 104891, and 105195) and the Finnish Environment Fund, JEF is supported by the MRC (Grant number G8802774), MJS by a grant from the British Heart Foundation, and MGM by an MRC Research Professorship.

Competing interests: none

REFERENCES

- 1 Stansfeld SA, Fuhrer R, Head J, et al. Work and psychiatric disorder in the Whitehall II Study. *J Psychosom Res* 1997;**43**:73–81.
- 2 Stansfeld SA, Rael EGS, Head J, et al. Social support and psychiatric sickness absence: a prospective study of British civil servants. *Psychol Med* 1997;**27**:35–48.
- 3 Stansfeld SA, Fuhrer R, Shipley MJ, et al. Work characteristics predict psychiatric disorder: prospective results from the Whitehall II Study. *Occup Environ Med* 1999;**56**:302–7.
- 4 Cheng Y, Kawachi I, Coakley EH, et al. Association between psychosocial work characteristics and health functioning in American women: prospective study. *BMJ* 2000;**320**:1432–6.
- 5 Escriba-Aguir V, Tenias-Burillo JM. Psychological well-being among hospital personnel: the role of family demands and psychosocial work environment. *Int Arch Occup Environ Health* 2004;**77**:401–8.
- 6 Kivimaki M, Elovainio M, Vahtera J. Workplace bullying and sickness absence in hospital staff. *Occup Environ Med* 2000;**57**:656–60.
- 7 Voss M, Floderus B, Diderichsen F. Physical, psychosocial, and organisational factors relative to sickness absence: a study based on Sweden Post. *Occup Environ Med* 2001;**58**:178–84.
- 8 Quine L. Workplace bullying in NHS community trust: staff questionnaire survey. *BMJ* 1999;**318**:228–32.
- 9 Varti MA. Consequences of workplace bullying with respect to the well-being of its targets and the observers of bullying. *Scand J Work Environ Health* 2001;**27**:63–9.
- 10 Mikkelsen EG, Einarsen S. Relationships between exposure to bullying at work and psychological and psychosomatic health complaints: the role of state negative affectivity and generalized self-efficacy. *Scand J Psychol* 2002;**43**:397–405.
- 11 Kivimaki M, Virtanen M, Varti MA, et al. Workplace bullying and the risk of cardiovascular disease and depression. *Occup Environ Med* 2003;**60**:779–83.
- 12 Elovainio M, Kivimaki M, Vahtera J. Organizational justice: evidence of a new psychosocial predictor of health. *Am J Public Health* 2002;**92**:105–8.
- 13 Miller DT. Disrespect and the experience of injustice. *Annu Rev Psychol* 2001;**52**:527–53.
- 14 Kivimaki M, Elovainio M, Vahtera J, et al. Organisational justice and health of employees: prospective cohort study. *Occup Environ Med* 2003;**60**:27–34.
- 15 Elovainio M, Kivimaki M, Vahtera J, et al. Sleeping problems and health behaviors as mediators between organizational justice and health. *Health Psychol* 2003;**22**:287–93.
- 16 Laaksonen M, Rahkonen O, Martikainen P, et al. Associations of psychosocial working conditions with self-rated general health and mental health among municipal employees. *Int Arch Occup Environ Health* 2006;**79**:205–12.
- 17 Sutinen R, Kivimaki M, Elovainio M, et al. Organizational fairness and psychological distress in hospital physicians. *Scand J Public Health* 2002;**30**:209–15.
- 18 Kivimaki M, Ferrie JE, Head J, et al. Organisational justice and change in justice as predictors of employee health: the Whitehall II study. *J Epidemiol Community Health* 2004;**58**:931–7.
- 19 Blaxter M. *Health and lifestyle*. London: Tavistock/Routledge, 1990.
- 20 Fylkesnes K, Forde OH. The Tromso Study: predictors of self-evaluated health—has society adopted the expanded health concept? *Soc Sci Med* 1991;**32**:141–6.
- 21 Smith AM, Shelley JM, Dennerstein L. Self-rated health: biological continuum or social discontinuity? *Soc Sci Med* 1994;**39**:77–83.
- 22 Singh-Manoux A, Martikainen P, Ferrie JE, et al. What does self-rated-health measure? Results from the British Whitehall II and French Gazel cohort studies. *J Epidemiol Community Health* 2006 (in press).
- 23 Goldberg DP. *The detection of psychiatric illness by questionnaire*. London: Oxford University Press, 1972.
- 24 Stansfeld SA, Marmot MG. Social class and minor psychiatric disorder in British Civil Servants: a validated screening survey using the General Health Questionnaire. *Psychol Med* 1992;**22**:739–49.
- 25 Lahelma E, Martikainen P, Rahkonen O, et al. Occupational class inequalities across key domains of health: results from the Helsinki Health Study. *Eur J Public Health* 2005;**15**:504–10.
- 26 Ferrie JE, Shipley MJ, Davey Smith G, et al. Change in health inequalities among British civil servants: the Whitehall II study. *J Epidemiol Community Health* 2001;**56**:922–6.
- 27 Gunnell D, Rasul F, Stansfeld SA, et al. Gender differences in self-reported minor mental disorder and its association with suicide. A 20-year follow-up of the Renfrew and Paisley cohort. *Soc Psychiatry Psychiatr Epidemiol* 2002;**37**:457–9.
- 28 Stansfeld SA, Fuhrer R, Shipley MJ, et al. Psychological distress as a risk factor for coronary heart disease in the Whitehall II Study. *Int J Epidemiol* 2002;**31**:248–55.
- 29 Rasul F, Stansfeld SA, Hart CL, et al. Psychological distress, physical illness, and risk of coronary heart disease. *J Epidemiol Community Health* 2005;**59**:140–5.
- 30 Laitinen-Krispijn S, Bijl RV. Mental disorders and employee sickness absence: the NEMESIS study. Netherlands Mental Health Survey and Incidence Study. *Soc Psychiatry Psychiatr Epidemiol* 2000;**35**:71–7.
- 31 Marmot MG, Davey Smith G, Stansfeld S, et al. Health inequalities among British civil servants: the Whitehall II study. *Lancet* 1991;**337**:1387–93.
- 32 Beksinska M, Yea L, Brunner E. *Whitehall II study: Manual for screening examination 1991–93*. London: Department of Epidemiology and Public Health, University College London, 1995.
- 33 Karasek R, Theorell T. *Healthy work: stress, productivity, and the reconstruction of working life*. New York: Basic Books, 1990.
- 34 Karasek R, Baker D, Marxer F, et al. Job decision latitude, job demands and cardiovascular disease: a prospective study of Swedish men. *Am J Public Health* 1981;**71**:694–705.
- 35 Karasek R. Job demands, job decision latitude, and mental strain: implications for job redesign. *Admin Sci Q* 1979;**24**:285–311.
- 36 North FM. Work and absence from work. Unpublished PhD thesis, University of London, 1990.
- 37 Bosma H, Marmot MG, Hemingway H, et al. Low job control and risk of coronary heart disease in the Whitehall II (prospective cohort) study. *BMJ* 1997;**314**:558–65.
- 38 Stansfeld SA, Bosma H, Hemingway H, et al. Psychosocial work characteristics and social support as predictors of SF36 health functioning: the Whitehall II study. *Psychosom Med* 1998;**60**:247–55.
- 39 Kuper H, Singh-Manoux A, Marmot M. When reciprocity fails: effort-reward imbalance in relation to coronary heart disease and health functioning within the Whitehall II study. *Occup Environ Med* 2002;**59**:777–84.
- 40 Nicholson A. Psychological distress as a predictor of coronary heart disease. Unpublished PhD thesis, University of London, 2003.
- 41 Rose GA, Blackburn H, Gillum RF, et al. *Cardiovascular survey methods*, 2nd edn. Geneva: WHO, 1982.
- 42 Kivimaki M, Elovainio M, Vahtera J, et al. Association between organizational inequity and incidence of psychiatric disorders in female employees. *Psychol Med* 2003;**33**:319–26.
- 43 Kivimaki M, Leino-Arjas P, Luukkonen R, et al. Work stress and risk of cardiovascular mortality: prospective cohort study of industrial employees. *BMJ* 2002;**325**:857.
- 44 Siegrist J. Adverse health effects of high-effort/low-reward conditions. *J Occup Health Psychol* 1996;**1**:27–41.
- 45 Siegrist J. Psychosocial work environment and health: new evidence. *J Epidemiol Community Health* 2004;**58**:888.
- 46 McAvoy BR, Murtagh J. Workplace bullying. *BMJ* 2003;**326**:776–7.
- 47 Ferrie JE, Shipley MJ, Marmot MG, et al. Health effects of anticipation of job change and non-employment: longitudinal data from the Whitehall II study. *BMJ* 1995;**311**:1264–9.
- 48 van den Bos K. Uncertainty management: the influence of uncertainty salience on reactions to perceived procedural fairness. *J Pers Soc Psychol* 2001;**80**:931–41.
- 49 Nunnally J. *Psychometric theory*. New York: McGraw-Hill, 1978.
- 50 Masterson SS, Lewis K, Goldman BM, et al. Integrating justice and social exchange: the differing effects of fairness procedures and treatment on work relationships. *Academy of Management Journal* 2000;**43**:738–48.
- 51 Elovainio M, Kivimaki M, Steen N, et al. Job decision latitude, organizational justice and health: multilevel covariance structure analysis. *Soc Sci Med* 2004;**58**:1659–69.
- 52 Mossholder KW, Bennett N, Martin CL. A multilevel analysis of procedural justice context. *Journal of Organisational Behaviour* 1998;**19**:131–41.
- 53 Schminke M, Cropanzano R, Rupp DE. Organization structure and fairness perceptions: the moderating effects of organizational level. *Organizational Behavior and Human Decision Processes* 2002;**89**:881–905.
- 54 Dailey RC, Kirk DJ. Distributive and procedural justice as antecedents of job dissatisfaction and intent to turnover. *Human Relations* 1992;**45**:305–17.

Commentary on

“Injustice at work and incidence of psychiatric morbidity: the Whitehall II study”.

Prepared for Occupational Environmental Medicine

Ichiro Kawachi
Harvard School of Public Health
677 Huntington Ave, Boston
MA 02115, USA.

ikawachi@hsph.harvard.edu

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in Occupational and Environmental Medicine editions and any other BMJ PGL products to exploit all subsidiary rights, as set out in our licence
<<http://OEM.bmjournals.com/misc/ifora/licenceform.shtml>>

Organizational justice has emerged in recent years as a determinant of workers' health, joining the growing list of other psychosocial aspects of the work environment, including job strain, effort-reward imbalance, and job insecurity. In a series of studies carried out mainly among Finnish workers, perceptions of organizational justice have been linked to poor self-rated health, minor psychiatric disorders, and sickness absences.^{1,2} In the current issue of the journal, Ferrie and colleagues provide an independent test of low organizational justice as a predictor of psychiatric morbidity within a well established cohort, the British Whitehall II study. What do these studies add to the literature on the psychosocial work environment, and do we have sufficient evidence to implicate organizational justice as a causal influence on workers' health?

Initial studies in this area were cross-sectional and involved self-reported outcomes, so that reverse causation and common method bias could not be ruled out. In a longitudinal follow-up of Finnish hospital workers, Kivimäki and colleagues² checked for the possibility of reverse causation by comparing the changes in perceptions of justice between initially healthy employees versus those with baseline health problems. Although the interaction term between time and baseline health was reported to be statistically non-significant, it was also evident that workers with health problems reported lower perceptions of justice compared to healthy co-workers at both the baseline and at the follow-up.² The new study by Ferrie *et al* examined the effects of change in relational justice over time in relation to the onset of psychiatric morbidity. A favorable change in perceptions of justice was associated with a reduction in psychiatric morbidity, whereas an adverse change increased the risk. These results bolster the case that reverse causation is not the major explanation for the observed associations. Nevertheless, if

health declines and changes in perceptions of justice are contemporaneous, it is difficult to completely rule out reverse causation, even with longitudinal change analysis. As Ferrie *et al* acknowledge, controlled experiments in the work setting would help, although it is not clear exactly what “treatment” should be designed to increase perceptions of justice.

Common method variance is a cause for concern when both the exposure and outcome variables are self-reported. Future studies of organizational justice and health would be strengthened by incorporating biomarkers and other endpoints that are not perceived or self-reported. Alternatively, common method variance could be addressed by aggregating individual responses to questions about organizational justice up to the work group or firm level. It makes theoretical sense to conceptualize and measure justice as an organization-level characteristic, as opposed to individual-level perceptions. Following this logic, investigators should focus on the contextual influence of organizational justice on workers’ health within a multi-level analytical framework, i.e. “exposure” to aggregated perceptions of justice assigned to individual workers nested within different workplaces. (A parallel argument could be made about investigating the health effects of other work environment characteristics, such as job strain and effort-reward imbalance).

A further noteworthy finding from existing studies of organizational justice and health is that the associations with endpoints tend to be attenuated (in several instances to statistical non-significance) after controlling for other psychosocial aspects of the work environment, including decision authority, effort-reward imbalance and social support. That is, the concept of organizational justice may be redundant to some degree with other

psychosocial aspects of work. As originally conceptualized by Moorman,³ organizational justice encompasses two domains: distributive justice and procedural justice. Because the definition of distributive justice (“the degree to which a worker believes that she is fairly rewarded in the basis of effort and performance”) essentially overlaps with Siegrist’s concept of effort-reward imbalance, subsequent research in the health field has focused on the procedural component of justice. According to Moorman, procedural justice, in turn, encompasses two dimensions: the existence of formal procedures in the workplace (i.e. the extent to which decision-making processes include input from affected parties, are fair and consistent, and provide useful feedback as well as the possibility of appeal), and interactional justice (the extent to which supervisors treat subordinates with respect, transparency, and fairness).³ In the public health field, the former dimension has been relabeled as “procedural justice”, while the latter is referred to as “relational justice”. However, as noted in an earlier commentary by Thoerell,⁴ the individual items in the procedural justice index overlap with the existing construct of decision authority, while the relational justice scale overlaps with the construct of supervisor support at work. Moreover, the correlation between the two components of organizational justice (procedural and relational) is only moderate (Pearson’s r about 0.3), while both constructs are also correlated to about the same degree with decision authority and workload.^{1,2} The new study by Ferrie *et al* was only able to examine the relational component of justice, although prior Finnish studies suggest that health (including psychiatric morbidity) is more strongly related to procedural justice than relational justice.^{1,2} Not surprisingly, because the 5-item scale of relational justice in the Whitehall II study was created by borrowing items from existing scales measuring effort-reward

imbalance and social support, the authors found that the biggest attenuations in odds ratios for psychiatric morbidity occurred after controlling for social support at work (among women), and effort-reward imbalance (among men). In summary, notwithstanding the finding that organizational justice is a statistically and empirically “independent” risk factor for workers’ health, greater clarity is called for in drawing out the theoretical distinctions as well as inter-relationships between justice and other established constructs in the psychosocial work environment.

A final point to note about studies of organizational justice is that employees with higher income levels have been reported to perceive lower levels of procedural justice,¹ so that differences in perceived organizational justice – in contrast to job control or effort-reward imbalance – is unlikely to explain the fundamental relationship between occupational status and workers’ health. Indeed, procedural justice has been reported to be a stronger predictor of sickness absence among high income hospital employees (rate ratio 1.17, 95% CI: 1.5 to 1.31) than among low income employees (RR = 0.95, 95% CI: 0.85 to 1.05), leading Kivimäki and colleagues to the somewhat counter-intuitive speculation that “procedural justice may have more salient meanings for members of highly ranked occupations close to management than for employees in lower ranking occupations”.^{2, p. 31} In the Whitehall II study, little association was found between relational justice scores and employment grade. Improving perceptions of procedural and relational justice is hence unlikely to contribute to reducing the social class gradient in psychiatric morbidity and other health problems. That is not to argue against attempting to intervene on perceptions of justice to improve workers’ health, but we must look

elsewhere to achieve “justice” (in its broader sense) in the distribution of work-related health outcomes.

References

1. **Elovainio M**, Kivimäki M, Vahtera J. Organizational justice: Evidence of a new psychosocial predictor of health. *Am J Public Health* 2002;**92**:105-108.
2. **Kimvimäki M**, Elovainio M, Vahtera J, Ferrie JE. Organisational justice and health of employees: prospective cohort study. *Occup Environ Med* 2003;**60**:27-33.
3. **Moorman RH**. Relationship between organizational justice and organizational citizenship behaviors: Do fairness perceptions influence employee citizenship? *J Appl Psychol* 1991;**76**:845-855.
4. **Theorell T**. Commentary. *Occup Environ Med* 2003;**60**:33-34.